

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Independent validation of the antiphospholipid score for the diagnosis of antiphospholipid syndrome.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/142601> since 2016-10-06T10:06:03Z

Published version:

DOI:10.1136/annrheumdis-2012-201985

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Sciascia S; Bertolaccini ML; Roccatello D; Khamashta MA. Independent validation of the antiphospholipid score for the diagnosis of antiphospholipid syndrome.. ANNALS OF THE RHEUMATIC DISEASES. 72 (1) pp: 142-143.

DOI: 10.1136/annrheumdis-2012-201985

The publisher's version is available at:

<http://ard.bmj.com/lookup/doi/10.1136/annrheumdis-2012-201985>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/142601>

Independent validation of the antiphospholipid score for the diagnosis of antiphospholipid syndrome

Savino Sciascia^{1,2}, Maria Laura Bertolaccini¹, Dario Roccatello², Munther A Khamashta^{1,3}

¹Lupus Research Unit, The Rayne Institute, Division of Women's Health, King's College, London, UK

²Centro di Ricerche di Immunopatologia e Documentazione su Malattie Rare (CMID), Università di Torino, Torino, Italy

³Louise Coote Lupus Unit, Guy's and St Thomas' NHS Foundation Trust, St Thomas' Hospital, London, UK

Antiphospholipid syndrome (APS) is a heterogeneous entity with a wide variation in clinical course and laboratory profile. It is accepted that the presence of antiphospholipid antibodies (aPL) confers a higher risk for both thrombosis and pregnancy morbidity, but quantifying such a risk is still a challenge. As a consequence, clinicians are unable to tailor the treatment according to the risk.

Recently, Otomo *et al*¹ developed and validated the so-called 'antiphospholipid score' (aPL-S) by testing multiple aPL and evaluating the aPL-S efficacy for the diagnosis of APS and predictive value for thrombosis. This score was shown to be a useful quantitative index for diagnosing APS and to be valuable as a predictive marker for thrombosis in autoimmune diseases.

In order to independently validate the aPL-S, we applied the proposed score system to a cohort of 211 consecutive patients who attended the Louise Coote Lupus Unit (St Thomas Hospital, London, UK). All the patients fulfilled the 1982 criteria for systemic lupus erythematosus (SLE).²

Overall, 81 patients fulfilled criteria for APS³ and 73 patients had a history of thrombosis (48 arterial, 41 venous thrombosis). Out of 144 women who had ever been pregnant, 41 had a history of miscarriages and 34 a history of fetal death. To validate the aPL-S, we adapted the proposed score using our inhouse cut-off values for aPL testing as previously reported or according to the current guidelines, as appropriate.³⁻⁸ aPL profile included anticardiolipin antibodies, lupus anticoagulant by partial thromboplastin time (aPTT – IL-test APTT-SP, Instrumentation Laboratory, Milan, Italy) and dilute Russell viper venom time, anti-β₂glycoprotein-I antibody, and antibodies to phosphatidylserine–prothrombin complex.

aPL-S was calculated for each patient by adding together the points corresponding to the risk factors as described.¹ Higher values of aPL-S were seen in patients who experienced thrombosis and/or pregnancy loss when compared with those without clinical events (median 17 (0–86) vs 4 (0–31), $p < 0.001$). When analysing clinical subgroups, patients who experienced thrombosis or pregnancy loss showed higher aPL-S compared with those without clinical events (median 18 (0–86) vs 4 (0–27), $p < 0.001$ for thrombosis; 7 (0–69) vs 3 (0–29), $p = 0.029$ for pregnancy loss) (figure 1). In our cohort, when the cut-off level for the aPL-S was defined as 30, as per the original study by Otomo *et al*,¹ the sensitivity and specificity of the aPL-S were 39% and 95%, respectively, compared with 37% and 96% shown in the Japanese cohort. The positive predictive value of an aPL-S ≥ 30 was 36%, whereas the negative predictive value was 91%.


We demonstrated that the aPL profile can be successfully quantified by the aPL-S in an independent cohort of SLE patients. The aPL-S correlated with a history of thrombosis or pregnancy loss in our cohort, suggesting that the aPL-S is a suitable quantitative marker of APS.

References

1. Otomo K, Atsumi T, Amengual O, *et al* Efficacy of the antiphospholipid score for the diagnosis of antiphospholipid syndrome and its predictive value for thrombotic events. *Arthritis Rheum* 2012;64:504–12.
2. Tan EM, Cohen AS, Fries JF, *et al* The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271–7.
3. Miyakis S, Lockshin MD, Atsumi T, *et al* International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295–306.

4. Brandt JT, Triplett DA, Alving B, *et al* Criteria for the diagnosis of lupus anticoagulants: an update. On behalf of the Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the ISTH. *Thromb Haemost*1995;74:1185–90.
5. Pengo V, Tripodi A, Reber G, *et al* Update of the guidelines for lupus anticoagulant detection. Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the International Society on Thrombosis and Haemostasis. *J Thromb Haemost* 2009;7:1737–40.
6. Harris EN, Gharavi AE, Boey ML, *et al* Anticardiolipin antibodies: detection by radioimmunoassay and association with thrombosis in systemic lupus erythematosus. *Lancet* 1983;2:1211–14.
7. Amengual O, Atsumi T, Khamashta MA, *et al*. Specificity of ELISA for antibody to beta 2-glycoprotein I in patients with antiphospholipid syndrome. *Br J Rheumatol*1996;35:1239–43.
8. Bertolaccini ML, Atsumi T, Koike T, *et al* Antiprothrombin antibodies detected in two different assay systems. Prevalence and clinical significance in systemic lupus erythematosus. *Thromb Haemost* 2005;93:289–97.

Figure 1



Antiphospholipid score (aPL-S) in systemic lupus erythematosus (SLE). The aPL-S was calculated according to Otomo *et al.*¹ Data are shown as box plots, where each box represents the 25th to 75th percentiles; lines inside the box represent the median. The whiskers represent the 95% CI. Higher values of aPL-S were seen in patients who experienced thrombosis and/or pregnancy loss (PL) when compared with those without clinical events ($p < 0.001$ by Mann–Whitney U test). When analysed separately, patients who experienced thrombosis or PL showed higher aPL-S when compared with those without clinical events.